



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

EXERCISES.

19

SHOW geometrically (without using the calculus) that the asymptote of the hyperbolic spiral, $r = a:\theta$, is parallel to the initial line and distant a above it.

[*L. G. Barbour.*]

20

A TUNNEL section, consisting of a rectangle surmounted by a semicircle, is required to accommodate a rectangle of breadth B and height H . Show how to determine its proportions so that its (1) area, (2) perimeter shall be least.

[*William M. Thornton.*]

21

IF THE central force on a body moving in a parabola, latus rectum $4m$, were to cease acting at the vertex and continue interrupted till the body had described an angle of 60° about the focus; determine the orbit it would afterwards describe.

[*William Hoover.*]

22

THE dome of the rotunda of the University of Virginia is spherical. The length of its meridian section is $85^f.2$ and the girth of its base is 214^f . From these data it is required to compute the radius and the surface of the dome.

[*William M. Thornton.*]

23

IF TWO parabolas P and P' cut each other at right angles at a point A on a third parabola P'' , all three parabolas having a common focus F , and if the tangent line drawn at A to P'' cut P and P' in B and C , then will one parabola that passes through B and C and has F for its focus cut P'' at right angles.

[*H. A. Newton.*]

24

The curves of the family

$$\left(\frac{C}{r}\right) = \cos(p + n\theta),$$

where n is a parameter, all pass through a fixed point and cut orthogonally the fixed curve

$$\left(\frac{r}{c}\right)^n = \cos n\theta,$$

provided $C^n = \frac{1}{2}c^n \cos p$.

[Generalization of 14.]

[*Alfred C. Lane.*]

25

AN elastic ring of radius a is placed gently on a smooth paraboloid of revolution, whose axis is vertical; find, by use of the principle of energy, the lowest position to which the ring will descend, and its position of static equilibrium. [R. D. Bohannon.]

NOTES.

3

MR. JOSEPH B. MOTT, of Worthington, Minn., sends, in connexion with an invalid deduction of the logarithmic series, several ingenious combinations for the computation of logarithms of primes. We note the following:—

$$\begin{aligned}\log 11 &= 1 + \frac{1}{2} \log 2 - 2 \log 3 + \log 7 \\ &\quad + M \left\{ \frac{1}{19601} + \frac{1}{3} \cdot \frac{1}{19601^3} + \frac{1}{5} \cdot \frac{1}{19601^5} + \dots \right\}. \\ \log 13 &= \frac{3}{2} \log 2 + \frac{3}{2} \log 11 - \log 3 - \frac{1}{2} \log 7 \\ &\quad - M \left\{ \frac{1}{21295} + \frac{1}{3} \cdot \frac{1}{21295^3} + \frac{1}{5} \cdot \frac{1}{21295^5} + \dots \right\}.\end{aligned}$$

Like results are given for $\log 17$ and $\log 19$; and the logarithms of these and other primes are computed with considerable facility to thirty-two places.

4

PERHAPS no modern geometer has fallen upon an easier and more rapid process for such computations than that indicated by Newton (*Epistola posterior ad Oldenburgium*, Oct. 24, 1676; *Opuscula* 1, 328). Newton computes for $x = 0,1$; $0,2$; $0,01$; $0,02$; $0,001$; $0,002$ the values of

$$\begin{aligned}\log \sqrt{\frac{1+x}{1-x}} &= x + \frac{1}{3}x^3 + \frac{1}{5}x^5 + \dots, \\ \log \sqrt{\frac{1}{1-x^2}} &= x^2 + \frac{1}{4}x^4 + \frac{1}{6}x^6 + \dots;\end{aligned}$$

whence by the aid of the simple interpolation formula

$$\log n = \log (n-x) + d \left(1 + \frac{x}{2n} + \frac{x^3}{12n^3} + \dots \right),$$